

Mepilex Border Sacrum How To

How To Use

Product description

Mepilex Border is an absorbent and self-adherent soft silicone dressing. It consists of:

- a soft silicone wound contact layer (Safetac).
- an absorbent pad in three layers: a polyurethane foam, a nonwoven layer and a fibre mixture with superabsorbent polyacrylate fibres.
- an outer film which is vapour permeable and waterproof.

Safetac Technology

Safetac is a patented soft silicone adhesive technology.

On the wound, Safetac assures non-sticking and minimal trauma on removal even in critical situations when the exudate starts to dry out.

On the skin, a dressing with Safetac has gentle but secure adhesion with no epidermal stripping or pain on removal.

Mode of action

Mepilex Border absorbs exudate, maintains a moist wound-healing environment and minimises the risk for maceration.

As Mepilex Border maintains a moist wound environment, supporting debridement, there might be an initial increase in the wound size. This is normal and to be expected. In case studies, dressings with Safetac technology have demonstrated to

normalise dry and scaly surrounding skin.

Due to the absorption and retention properties Mepilex Border is suitable for use under compression bandaging. Mepilex Border can be used in combination with gels.

Intended use

Mepilex Border is designed for a wide range of exuding wounds such as pressure ulcers, leg and foot ulcers and traumatic wounds e.g skin tears and surgical wounds. Mepilex Border can also be used on dry/necrotic wounds in combination with gels.

Instructions for use

1. Cleanse the wound in accordance with normal procedures.
2. Dry the surrounding skin thoroughly.
3. Remove the release films.
4. Apply the adherent side to the wound. Do not stretch.
5. On dressing removal gently lift one corner and slowly peel back the dressing.
6. If extra fixation is needed – fixate only on the borders of the product.

Frequency of change

Mepilex Border may be left in place for several days depending on the condition of the wound and the surrounding skin, or as indicated by accepted clinical practice.

Mepilex Border should be changed when exudate is present at the pad edges.

A change in dressing regimen can result in an initial increased level of exudate which temporarily may require an increased change frequency.

Precautions

In case of signs of clinical infection, consult a health care professional for adequate infection treatment. Do not use Mepilex Border together with oxidising agents such as hypochlorite solutions or hydrogen peroxide. Mepilex Border should be stored in dry conditions below 35°C (95F) and be protected from direct sunlight. Do not use if inner package is damaged. Sterility is guaranteed if inner package is intact. Do not resterilise.

Mepilex and Safetac are registered trademarks of Mölnlycke Health Care AB.